

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets

(11) Publication number:

0 236 930
A1

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: 87103077.1

(22) Date of filing: 04.03.87

(51) Int. Cl.³: **C 07 D 275/06**
C 07 D 417/12, C 07 D 471/0-4
A 61 K 31/425, A 61 K 31/53-5
A 61 K 31/54, A 61 K 31/40
A 61 K 31/495, A 61 K 31/43-5
/(C07D471/04, 221:00, 209:00)


(30) Priority: 05.03.86 US 836276

(43) Date of publication of application:
16.09.87 Bulletin 87/38(84) Designated Contracting States:
AT BE CH DE ES FR GB GR IT LI LU NL SE(71) Applicant: MERRELL DOW PHARMACEUTICALS INC.
2110 E. Galbraith Road
Cincinnati Ohio 45215(US)(72) Inventor: Hibert, Marcel
6 Rue Richard Wagner
F-67000 Strasbourg(FR)(72) Inventor: Gittos, Maurice W.
16 Rue André Malraux
F-67115 Plobsheim(FR)(74) Representative: Macchetta, Francesco et al,
Gruppo Lepetit S.p.A. Patent and Trademark Department
34, Via Roberto Lepetit
I-21040 Gerenzano (Varese)(IT)

(54) Aromatic 2-aminoalkyl-1,2-benzisothiazol-3(2H)one-1,1-dioxide derivatives.


(57) Aromatic 2-amino-alkyl-1,2-benzisothiazol-3(2H)one-1,1-dioxide derivatives having the general formula

thereof, a process for the preparation of these compounds and discloses their use as anxiolytic and antihypertensive agents.



(1)

wherein X is selected from the group consisting of



n is an integer of from 2 to 5; A and B are each oxygen sulfur or NR' where R' represents hydrogen or methyl; R₁ is hydrogen, hydroxy or methoxy; R₂ is hydrogen or methyl; and the pharmaceutically acceptable acid addition salts

AROMATIC 2-AMINOALKYL-1,2-BENZOISOTHIAZOL-3(2H)ONE-1,1-DIOXIDE
DERIVATIVES

FIELD OF THE INVENTION

5 This invention relates to certain aromatic 2-aminoalkyl-
1,2-benzothiazol-3(2H)one-1,1-dioxide derivatives and their
use as anxiolytic agents and antihypertensive agents.

BACKGROUND OF THE INVENTION

10 Anxiety has been defined as an apprehension or concern
regarding some future event. Most, if not all, people
occasionally suffer some symptoms of anxiety in response to
appropriate stimuli. In some individuals, these feelings of
anxiety or panic in response to the everyday pressures of life
can be overwhelming, rendering the individual an unproductive
15 member of society. Whereas individual group counseling
represents the preferred primary mode of therapy, the use of
chemotherapeutic agents has proven to be a useful adjunct in

the treatment of anxiety, thereby enabling a seriously afflicted individual to regain productive status while undergoing concurrent psychotherapy.


Compounds of the class of benzodiazepines are currently the therapeutic agents of choice in the treatment of anxiety. In particular, chlordiazepoxide, diazepam and oxazepam are commonly used. This class of compounds has a great potential for misuse, particularly among the class of patients undergoing therapy. Moreover, the benzodiazepines generally possess undesired sedative effects and process detracting interactions with other drugs, including for example, alcohol.

Applicants have now discovered a class of novel aromatic 2-aminoalkyl-1,2-benzothiazol-3(2H)one-1,1-dioxide antianxiety agents that are generally free from the undesirable effects of the benzodiazepines. The compounds disclosed herein, when practised in accordance with the teachings of this invention help to alleviate such symptoms as excessive fear, worry, restlessness, tension, stress, neurotic depression and are useful in the relief of some personality disorders. Additionally, the compounds of this invention are effective antihypertensive agents which have the effect of lowering blood pressure in patients in need thereof.

SUMMARY OF THE INVENTION


This invention is directed to a class of aromatic 2-amino-alkyl-1,2-benzothiazol-3(2H)one-1,1-dioxide derivatives having the general formula

5





(1)

10 wherein X is selected from the group consisting of




15






5



10



15

n is an integer of from 2 to 5; A and B are each oxygen, sulfur or NR' where R' represents hydrogen or methyl; R₁ is hydrogen, hydroxy or methoxy; R₂ is hydrogen or methyl; and the pharmaceutically acceptable acid addition salts thereof.


This invention also discloses a process for the preparation of these compounds and discloses their use as anxiolytic and antihypertensive agents.

DETAILED DESCRIPTION OF THE INVENTION

As used herein the term "pharmaceutically acceptable acid addition salts" is intended to apply to any non-toxic organic or inorganic acid addition salts of the base compounds represented by Formula I. Illustrative inorganic acids which form suitable salts include hydrochloric, hydrobromic, sulphuric and phosphoric acid and acid metal salts such as sodium monohydrogen orthophosphate and potassium hydrogen sulfate. Illustrative organic acids which form suitable salts include the mono, di and tricarboxylic acids. Illustrative of such acids are, for example, acetic, glycolic, lactic, pyruvic, malonic, succinic, glutaric, fumaric, malic, tartaric, citric, ascorbic, maleic, hydroxymaleic, benzoic, p-hydroxybenzoic, phenylacetic, cinnamic, salicylic, 2-phenoxybenzoic acids, and sulfonic acids such as methane-sulfonic acid or 2-hydroxyethanesulfonic acid. Either the mono or the di-acid salts can be formed, and such salts can exist in either a hydrated or a substantially anhydrous form. In general, the acid addition salts of these compounds are crystalline materials which are soluble in water and in various hydrophilic organic solvents. Additionally, in comparison to their free base forms, such salts generally demonstrate higher melting points and an increased chemical stability.


Specific subclasses of compounds that fall within the scope of the present invention are illustrated as follows:

2-[ω -[(2,3-dihydro-1,4-benzodioxin-2-yl)methylamino]alkyl]-1,2-benzoisothiazol-3(2H)one-1,1-dioxides



(1a)


2-[ω -[(2,3-dihydro-1,4-benzoxathiin-3-yl)methylamino]alkyl]-1,2-benzoisothiazol-3(2H)one-1,1-dioxides



(1b)

2-[ω-[(2,3-dihydro-1,4-benzodithiin-2-yl)methylamino]alkyl]-
1,2-benzoisothiazol-3(2H)one-1,1-dioxides


5



(1c)

10

2-[ω-[(2,3-dihydro-1,4-benzoxazin-3-yl)methylamino]alkyl]-
1,2-benzoisothiazol-3(2H)one-1,1-dioxides




15

(1d)

2-[ω-[(2,3-dihydroquinazolin-2-yl)methylamino]alkyl]-
1,2-benzoisothiazol-3(2H)one-1,1-dioxides


20



(1e)

2-[ω-[(2,3-dihydronaphtho[1,2-b][1,4]dioxin-2-yl)methylamino]alkyl]-1,2-benzisothiazol-3(2H)one-1,1-dioxides **0236930**

5




(Compounds of this subclass can be substituted either at the 2- or 3-position of the dioxin ring.)

10

(1f)

2-[ω-[substituted (indol-3-yl)ethylamino]alkyl]-1,2-benzisothiazol-3(2H)one-1,1-dioxides


15



(1g)

2-[ω-[(substituted)1,2,3,4-tetrahydro-β-carboliny]alkyl]-
1,2-benzisothiazol-3(2H)one-1,1-dioxides


5



(1h)

2-[ω-[(substituted)-tetralin-2-amino]alkyl]1,2-benzisothiazol-3(2H)one-1,1-dioxides

10



15

(1i)

A preferred subclass of this invention consists of those compounds of formula (1h) wherein the alkyl chain is terminally substituted with a (substituted)-1,2,3,4-tetrahydro-β-carboline ring system.

20


0236930

A more preferred subclass of this invention relates to those compounds of formula (1f) wherein the alkyl chain is terminally substituted by the 2,3-dihydronaphtho[1,2-b][1,4]-dioxin-2-yl or 3-yl ring system.

5 The most preferred compounds of this invention relate to those compounds of formula (1a) wherein the alkyl chain is terminally substituted by the 2,3-dihydro-1,4-benzodioxin-2-yl ring system.

10 The alkylene group depicted above $-(CH_2)_n-$, can be considered as a connecting bridge which separates the two terminal heterocyclic ring systems. As indicated, the symbol n can represent an integer of from 2 to 5. Those alkylene groups in which n is from 2 to 4 represent the preferred groups of this invention.

15 The aromatic 2-aminoalkyl-1,2-benzoisothiazol-3(2H)one-1,1-dioxide derivatives of formula (1) can be prepared in an analogous manner by standard techniques known to those skilled in the art. Thus, the compounds of this invention can be prepared via a condensation of the appropriate nucleophilic
20 amine of formula (2) with an N-alkyl-1,2-benzoisothiazol-3(2H)one-1,1-dioxide substrate of formula (3) as indicated below



wherein X and n are as defined in formula (1) and the symbol (L) represents a suitable leaving group, such as chlorine, bromine, iodine, a mesylate or tosylate.

Such a nucleophilic condensation is preferably conducted by reacting approximately equimolar amounts of the nucleophile (2) with the substrate (3), for a period of from about 1 hour to 24 hours depending upon the particular reactants employed. The reaction temperature can range from about 25°C to 140°C. Preferably the reaction is conducted at a temperature ranging from 60°C to 125°C.


Additionally, the reaction is preferably conducted in the presence of a tertiary organic base, such as a trialkylamine or pyridine, or in the presence of an inorganic base such as potassium carbonate.

Inasmuch as the reactants employed are typically crystalline materials, the use of solvents is preferred. Suitable solvents include any non-reactive solvent, preferably those having a boiling point in the range of from 60°C to 150°C. Thus, for example, solvents such as petroleum ethers; chlorinated


hydrocarbons such as carbon tetrachloride, ethylene chloride, methylene chloride or chloroform; chlorinated aromatic compounds such as 1,2,4-trichlorobenzene, or o-dichlorobenzene; carbon disulfide; ethereal solvents such as diethylether, tetrahydrofuran or p-dioxane; aromatic solvents, such as benzene, toluene or xylene; or an alcoholic solvent such as ethanol, can be suitably employed. Especially preferred solvents are those which are known to promote nucleophilic reactions, such as dimethylsulfoxide and dimethylformamide.

The products of formula (1) can be isolated utilizing appropriate techniques available to those skilled in the art. Thus, for example, the reaction mixture can be filtered to remove solid materials, and the filtrate subsequently evaporated to recover the active ingredient. The compounds of formula (1) can be further purified via recrystallization or by forming and utilizing their picric or oxalic acid complexes.

The nucleophilic primary amines indicated by formula (2) are compounds which are either commercially available or which have been previously described in the literature. Alternatively, the primary amines, designated by the symbol X, wherein R_2 is hydrogen, can be readily prepared via a reduction of the corresponding cyano derivatives. Thus, for example, compounds of formula (2), wherein X represents the radical




in which A and B are as defined in formula (1) above, are prepared by a reduction of the corresponding cyano derivative
 5 (4) shown below



(4)

Such a reduction can be accomplished utilizing a variety of
 10 reagent systems, as for example, catalytic reductions employing hydrogen gas with a catalytic metal such as palladium on charcoal, Raney nickel, platinum, rhodium, ruthenium or platinum oxide. In addition, reagents such as diborane, sodium borohydride, dissolving metal reductions
 15 utilizing lithium, sodium, potassium, calcium, zinc, magnesium, tin or iron in liquid ammonia or a low-molecular weight aliphatic amine or sodium, aluminum or zinc amalgam, zinc, tin or iron in a hydroxylic solvent or in the presence of an aqueous mineral acid, or lithium aluminum hydride can be
 20 favorably employed.


The formula (2) nucleophiles wherein X represents the radical




5 can be prepared by reacting the corresponding cyano compounds with from 1 to 2 molar equivalents of lithium aluminum hydride in an appropriate solvent. Preferably, about 1.5 molar equivalents of hydride are employed. The reaction is allowed to proceed for a period of time ranging from about 30 minutes
10 to about 24 hours. Preferably a time period of from about 1 to 5 hours is employed, depending upon the particular reactants, the solvent and temperature utilized. Suitable temperatures range from -78°C to 60°C , preferably about 20°C . Suitable solvents include ethereal solvents such as diethyl ether,
15 tetrahydrofuran (THF), p-dioxane, 1,2-dimethoxyethane (DME), diglyme or an aromatic solvent such as benzene, toluene or xylene.

In general, the secondary nucleophilic amines of formula (2), wherein X represents the radicals


20





and R_2 is other than hydrogen, can be prepared by the direct alkylation of the corresponding primary amine with a suitable alkyl halide, tosylate or mesylate in an appropriate solvent, such as acetonitrile, in the presence of at least one equivalent or an excess of an organic or inorganic base, such as potassium carbonate.

Many of the cyano derivatives of formula (4) are known compounds. Alternatively, these compounds can be prepared in an analogous manner as described below. Thus, for example, utilizing the same example as previously employed the compound



(5)


wherein A and B are as described above, is reacted with 2-bromo or preferably 2-chloroacrylonitrile. Approximately, equimolar amounts of the formula (5) compound and the

haloacrylonitrile are mixed with 2 or more molar equivalents of a base, such as potassium carbonate, in an appropriate solvent. The reaction is allowed to proceed at a temperature ranging from 0°C to the boiling point of the reaction mixture, for a period ranging from about 1 to 24 hours. Suitable solvents employed include dimethylformamide; dimethylsulfoxide; acetone; chlorinated hydrocarbons, such as carbon tetrachloride, chloroform or methylene chloride; ethereal solvents, such as diethylether, tetrahydrofuran (THF) or diglyme; aromatic solvents, such as benzene, toluene or xylene; or alcoholic solvents, such as methanol or ethanol.

Where the symbols A and B represent different atoms, a mixture of products will be obtained. These mixtures can be readily separated and purified by methods commonly known to those skilled in the art, such as by chromatography on silica gel or fractional recrystallization. Furthermore, when the R₁ or R₂ substituents of a compound of formula (1) are hydroxy, the hydroxyl group must be protected prior to undergoing the above described 2-bromo- or 2-chloroacrylonitrile condensation reaction. Suitable protecting groups include the benzyl or methyl groups. These protecting groups are subsequently removed to form the corresponding nucleophilic amines represented by formula (2). The removal of such protecting groups can be by any suitable means generally known to the art, such as the

catalytic reduction of the benzyl group, treatment with an acid such as hydrobromic acid, or treatment with boron tri-bromide.

The compounds of formula (3) are essentially N-alkyl derivatives of saccharin. The leaving groups (L) for compounds of formula (3) can represent any group known to those skilled in the art, such as a tosylate (OTS) or mesylate (OMS), an iodide, bromide or chloride, or hydroxyl group. Formula (3) substrates wherein L is a bromide or an iodide can be prepared by treating the corresponding dihaloalkane with one equivalent of the sodium salt of saccharin in an appropriate solvent for about one hour at a temperature of about 100°C, in accordance with the following reaction scheme.



wherein the symbol Hal represents chloro, bromo or iodo.

Suitable solvents include dimethylformamide; dimethylsulfoxide; acetone; aromatic solvents, such as benzene, toluene or xylene; or an ethereal solvent, such as diethyl ether, tetrahydrofuran (THF) or 1,2-dimethoxyethane (DME).

The compounds of formula (1) possess useful antianxiety and antihypertensive properties. Anxiolytic properties are suggested using 5-HT_{1A} in vitro receptor binding studies, see Middlemiss et al., Eur. J. Pharmacol., 90, 151-3 (1983) and Glaser et al., Arch. Pharmacol., 329, 211-215 (1985). The antihypertensive effects of the compounds described herein can be determined both in the anesthetized normotensive rat and/or in the conscious spontaneously hypertensive rat in accordance with the procedure of Fozard, J. Cardiovascular Pharm., 4, 829-838 (1982).

The compounds of this invention can be administered either orally, subcutaneously, intravenously, intramuscularly, intraperitoneally or rectally. The preferred route of administration is oral. The amount of compound to be administered can be any effective amount, and will vary depending upon the patient, the mode of administration and the severity of the anxiety to be treated. Repetitive daily administration of the compounds may be desirable, and will vary depending upon the patient's condition and the mode of administration.

For oral administration, an anxiolytic or antihypertensive effective amount of a formula (1) compound can range from 0.005 to 10 mg/kg of patient body weight per day, preferably from 0.05 to 5 mg/kg of patient body weight per day. The preferred antianxiety dose of the compounds of formula (1a) is about 0.1 mg/kg of patient body weight per day. Pharmaceutical compositions in unit dose form can contain from 1 to 50 mg of active ingredient and can be taken one or more times per day.


For parenteral administration, an anxiolytic or antihypertensive effective amount of a formula (1) compound is from about 0.005 to 10 mg/kg of patient body weight per day, preferably from about 0.05 to 5 mg/kg of patient body weight per day. A parenteral composition in unit dose form can contain from 0.1 mg to 10 mg of active ingredient and can be taken one or more times daily.

For oral administration the compounds can be formulated into solid or liquid preparations such as capsules, pills, tablets, lozenges, melts, powders, solutions, suspensions or emulsions. Solid dosage unit forms generally employed include capsules or tablets. Capsules can be of the ordinary gelatin type which contain additional excipients such as, surfactants, lubricants and inert fillers such as lactose, sucrose and cornstarch. Additionally, the compounds of formula (1) can be tableted with conventional tablet bases such as lactose, sucrose, and cornstarch in combination with binders, such as acacia,

cornstarch or gelatin, disintegrating agents such as potato starch or alginic acid, and lubricants such as stearic acid or magnesium stearate.

For parenteral administration the compounds may be administered as injectable dosages of a solution or a suspension of the compound in a physiologically acceptable diluent with or without a pharmaceutical carrier. Suitable diluents or carriers include sterile liquids such as water or oils, with or without the addition of surfactants or other pharmaceutically acceptable adjuvants. Illustrative of various oils that can be employed in the practice of this invention are those of petroleum, animal, vegetable, or synthetic origin, for example, peanut oil, soybean oil, and mineral oil. In general, water, saline, aqueous dextrose and related sugar solutions, ethanol and glycols such as propylene glycol or polyethylene glycol are preferred liquid carriers, particularly for injectable solution.

The following examples illustrate the preparation of representative compounds employed in the practice of this invention, but are not intended to limit the invention in any way thereto.

EXAMPLE I

2-[4-[(2,3-Dihydro-1,4-benzodioxin-2-yl)methylamino]butyl]-1,2-benzoisothiazol-3(2H)one-1,1-dioxide

N-(4-Bromobutyl)saccharin

Saccharin (9.35g, 50mM) in 80ml of N,N-dimethyl formamide (DMF) is added to 55% sodium hydride (2.18g, 50mM) in 20ml of dry DMF. After 15 minutes, 1,4-dibromobutane (25ml, 200mM) is added rapidly, and the mixture is warmed for 1 hour at 100°C with stirring. The reaction mixture is cooled, filtered and the DMF is evaporated. The crude product is dissolved in methylene chloride and the solution is filtered, dried and evaporated, yielding a yellow oil which is purified by chromatography on silica gel (CH₂Cl₂/MeOH, 98/2) to afford 13.4g (84%) of a colorless oil. This oil, when crystallized from isopropanol, yields N-(4-bromobutyl)saccharin having a melting point of 72°C.

NMR CDCl_3/TMS 60 MHz 8.20-7.85 (m, 4 H Ar), 4.00-3.30 (m, 4 H), 2.40-1.80 (m, 4H).


2-[4-[(2,3-Dihydro-1,4-benzodioxin-2-yl)methylamino]butyl]-1,2-benzoisothiazol-3(2H)one-1,1-dioxide hydrochloride

5 2-Aminomethyl-benzodioxan[1,4] (0.965g, 5.48mM), potassium carbonate (3g) and N-(4-bromobutyl)-saccharin (1.43g, 4.5mM), prepared as above, are mixed in N,N-dimethylformamide (DMF, 10ml). The mixture is stirred at 100°C overnight, cooled, filtered and evaporated to dryness. The residue is dissolved
0 in ethyl acetate, washed with water and acidified with dilute hydrochloric acid. The oil which separates is washed with ether, 5% HCl, water, dissolved in methanol, dried and evaporated to dryness, affording 1.3g of a crude yellow solid. This material is converted to the free base and flash-
.5 chromatographed on silica ($\text{CH}_2\text{CO}_2/\text{MeOH}$ 97/3), yielding 0.91g of the desired compound. The hydrochloride salt is recrystallized from an isopropanol/ethyl acetate solution to yield the title compound having a melting point of 210°C.

Following essentially the same procedure but substituting
20 N-(2-bromoethyl)saccharin for the N-(4-bromobutyl)saccharin above, the compound 2-[2-[(2,3-dihydro-1,4-benzodioxin-2-yl)-methylamino]ethyl]-1,2-benzoisothiazol-3(2H)one-1,1-dioxide is obtained, having a melting point of 181°C.

EXAMPLE II


0236930



5


2-[4-(2-[1,2,3,4]-tetrahydro- β -carbolinyl)butyl]-1,2-benzo-
isothiazol-3(2H)one-1,1-dioxide

Following essentially the same procedure as in Example I,
but substituting [1,2,3,4]-tetrahydro- β -carboline for
10 2-aminomethyl-benzodioxan[1,4], the title compound was
prepared and crystallized as a hydrochloride, having a melting
point of 272°C.

EXAMPLE III

2-[4-[8-Methoxytetralin-2-amino]butyl]1,2-benzisothiazol-
3(2H)one-1,1-dioxide


Following essentially the same procedure as in Example I,
10 but substituting 8-methoxy-2-aminotetralin for 2-aminomethyl-
benzodioxan[1,4], the title compound was prepared and
crystallized as a hydrochloride, having a melting point of
233°C.

EXAMPLE IV

2-[4-[5-Methoxy-indol-3-yl)ethylamino]butyl]1,2-benzothiazol-3(2H)one-1,1-dioxide

Following essentially the same procedure as in Example I, but substituting 5-methoxy-tryptamine for 2-aminomethyl-benzodioxan[1,4], the title compound was prepared and crystallized as a hydrochloride, which decomposes prior to melting.

¹H NMR (free base, CDCl₃+CD₃OD, 360 MHz, ppm): 8.4 (1H,s), 8.0-7.7 (4H,m), 7.2-6.75 (4H,m), 3.8 (3H,s), 3.8-3.6 (2H,m), 3.15 (4H,m), 2.9 (2H,t), 1.85 (4H,m).

EXAMPLE V

2-[4-[(2,3-Dihydro-1,4-benzoxazin-3-yl)methylamino]butyl]-1,2-benzoisothiazol-3(2H)one-1,1-dioxide

Following essentially the same procedure as in Example I,
10 but substituting 2-aminomethylbenzoxazin for 2-aminomethyl-
benzodioxan[1,4], the title compound was prepared and
crystallized as a hydrochloride, having a melting point of
209°C.

EXAMPLE VIIn vitro Determination of Anxiolytic Propertiesvia 5-HT_{1A} Binding

5 Radioligand binding studies of the 5-HT_{1A} recognition sites
were carried out as follows: male normotensive Sprague-Dawley
rat frontal cortex was dissected, frozen in liquid nitrogen
and stored at -20°C until needed. Tissues from 4-8 rats were
pooled and homogenised in 70 vol Tris-HCl buffer (50mM,
pH 7.7), using a kinematica Polytron (setting 2/3 max speed,
10 20 sec). The homogenate was centrifuged (36500 x g for
10 min), the pellet re-homogenised in the same volume of
buffer and the process repeated two more times. Between the
second and third centrifugations the tissue homogenate was
incubated at 37°C for 10 min. The final pellet was suspended
15 in the same volume of Tris buffer containing 10 M pargyline,
5.7mM CaCl₂ and 0.1% ascorbic acid. This suspension was
incubated for 10 min at 37°C and then stored on ice until used
in the binding assay.

20 Tissue homogenate (0.7ml), radioactive ligand (0.1ml) and
the appropriate concentration of test compound (0.1ml),
together with buffer to a final volume of 1ml were incubated
at 37°C for 15 min. Incubations were terminated by rapid
filtration through Whatman GF/B filters followed by three 5ml

washes with ice-cold Tris-HCl buffer (50mM, pH 7.0). Radio-activity was measured after extraction into Aquasol-Z (NEN) at an efficiency of 45-50%. The radioligand used to label the 5-HT_{1A} recognition sites and its concentration is [³H]-8-hydroxy-2-(di-n-propylamino)-tetralin, ([³H]-8-OH-DPAT), 1mM.

Following essentially the above procedure, the following compounds were tested. Results are expressed as pIC₅₀ (log₁₀ concentration of test compound which inhibits specific binding by 50%), and represent the means \pm S.E.M. of 3 independent experiments.

Test Compound	5-HT _{1A} Binding Affinity Rat Brain Cortex	* Fold Increase
Buspirone	7.52 \pm 0.10	1
Example I(A)	9.34 \pm 0.03	66
Example II	8.16 \pm 0.07	4.4
Example III	8.52 \pm 0.14	10
Example V	8.03 \pm 0.15	3.2

* Antilog of the difference of the test compound compared to buspirone

As indicated in the above data, the compounds tested represent a 3 to 60 fold increase in potency, based upon 5HT_{1A} binding studies, when compared to buspirone.

EXAMPLE VIIDetermination of Antihypertensive properties

Male, normotensive Sprague-Dawley rats weighing 250-350g (supplied by Charles River, France) were anesthetized with pentobarbitone sodium, 60mg/kg, i.p., plus 15mg/kg s.c. Blood pressure was recorded either from the left common carotid artery or from a femoral artery by means of a Statham pressure transducer (Type P23 AA) and heart rate was recorded from the electrocardiogram using a Beckman cardiometer coupler (Type 9757 B). Records were displayed on a Beckman dynograph (Type R). A femoral vein was cannulated for the intravenous injection of drugs. After completion of all operative procedures, heparin 500 U was injected intravenously.

The compounds of Example I, 2-[4-[(2,3-dihydro-1,4-benzodioxine-2-yl)methylamino]butyl]-1,2-benzisothiazol-3(2H)one-1,1-dioxide was administered s.c. as a saline solution in a volume of 1ml/kg body weight. The following changes in blood pressure (B.P.) and heart rate (H.R.) were observed.

20	Dose	Δ BP (mmHg)	Δ HR (beats/min)
	25 μ g/kg	-35	-80
25	500 μ g/kg	-50	-90


In a separate experiment arterial blood pressure and heart rate were measured directly in conscious male, spontaneously hypertensive rats weighing 250-350g (Charles River, France). The same test compound was administered, either s.c. or orally, as a saline solution of 1ml/kg body weight in the doses shown below.

	Route and Dose	Δ BP (mmHg)	Δ HR (beats/min)
10	100μg/kg s.c.	-20	+80
	200μg/kg s.c.	-65	+100
15	500μg/kg s.c.	-75	+120
	1mg/kg per os.	-25	+100
20	5mg/kg per os.	-45	+100




In short, under the conditions tested, it can be said that the compound, 2-[4-[(2,3-dihydro-1,4-benzodioxin-2-yl)methyl-amino]butyl]1,2-benzoisothiazol-3(2H)one-1,1-dioxide effectively lowers the blood pressure and heart rate in anesthetized normotensive rats, and lowers the blood pressure with reflex tachycardia in conscious spontaneously hypertensive rats.



CLAIMS FOR THE CONTRACTING STATES: BE, FR, DE, GB, IT,
LU, NL, SE, CH.

1. An aromatic 2-aminoalkyl-1,2-benzothiazol-3(2H)one-1,1-dioxide derivative having the formula




wherein n is an integer of from 2 to 5; X is selected from the group consisting of





A and B are each oxygen, sulfur or NR' where R' is hydrogen or methyl; R₁ is hydrogen, hydroxy or methoxy; R₂ is hydrogen or methyl; and the pharmaceutically acceptable acid addition salts thereof.


2. A compound according to Claim 1 wherein X is CH₂NH-; A and B are both oxygen, n is an integer of from 2 to 4, and the pharmaceutically acceptable acid addition salts thereof.

3. A compound according to Claim 2 which is 2-[4-[(2,3-dihydro-1,4-benzodioxin-2-yl)methylamino]butyl]-1,2-benzothiazol-3(2H)one-1,1-dioxide or the pharmaceutically acceptable salts thereof.

0236930

4. A compound according to Claim 2 which is 2-[2-[(2,3-dihydro-1,4-benzodioxin-2-yl)methylamino]ethyl]-1,2-benzothiazol-3(2H)one-1,1-dioxide and the pharmaceutically acceptable salts thereof.


5. A compound according to Claim 1 wherein X is



4, and the pharmaceutically acceptable salts thereof.

6. A compound according to Claim 5 which is 2-[4[(2,3-dihydronaphtho[1,2-b][1,4]dioxin-3-yl)methylamino]butyl]-1,2-benzothiazol-3(2H)one-1,1-dioxide and the pharmaceutically acceptable salts thereof.

7. A compound according to Claim 1 wherein X is





n is an integer of from 2 to 4, and the pharmaceutically acceptable salts thereof.


8. A compound according to Claim 7 which is 2-[4-(1,2,3,4-tetrahydro- β -carboliny1)butyl]-1,2-benzisothiazol-3(2H)one-1,1-dioxide and the pharmaceutically acceptable salts thereof.

9. A process for the preparation of an aromatic 2-aminoalkyl-1,2-benzisothiazol-3(2H)one-1,1-dioxide derivative of Claim 1, or a pharmaceutical salt thereof, which comprises reacting a nucleophilic amine having the formula:




wherein X is selected from the group consisting of






5




or



- 10 wherein the symbols A, B, R₁ and R₂ are as defined in Claim 1;
with an N-alkyl-1,2-benzothiazol-3(2H)one-1,1-dioxide
substrate having the formula

15




wherein n is as defined in Claim 1, and L is a leaving group;
for a period of from 1 to 24 hours and a temperature of from
25°C to 140°C; and isolating the desired product therefrom.




10. A compound of claim 1 to 8 for use as a medicine.
- 5 11. Use of a compound of claim 1 to 8 for the preparation of a medicament for use as an ansiolytic agent.
- 10 12. Use of a compound of claim 1 to 8 for the preparation of a medicament for use as an antihypertensive agent.
- 15 13. A pharmaceutical composition comprising a compound of claim 1 to 8 in admixture with a pharmaceutically acceptable carrier.



CLAIMS FOR THE CONTRACTING STATES: AT, ES, GR.

- 5 1. A process for the preparation of 2-aminoalkyl-1,2-benzisothiazol-3(2H)one-1,1-dioxide derivative having the formula (1)



- 10 wherein n is an integer of from 2 to 5; X is selected from the group consisting of









A and B are each oxygen, sulfur or NR' where R' is hydrogen or methyl; R₁ is hydrogen, hydroxy or methoxy; R₂ is hydrogen or methyl; or a pharmaceutical salt thereof, which comprises reacting a nucleophilic amine having the formula:




wherein X is selected from the group consisting of






, or



wherein the symbols A, B, R₁ and R₂ are as defined above; with an N-alkyl-1,2-benzothiazol-3(2H)one-1,2-dioxide substrate having the formula



wherein n is as defined above, and L is a leaving group; for a period of from 1 to 24 hours and a temperature of from 25°C to 140°C; and isolating the desired product therefrom.

2. A process according to claim 1 for preparing a

compound of formula (1) wherein X is


A and B are both oxygen, n is an integer of from 2 to 4, and the pharmaceutically acceptable acid addition salts thereof.

3. A process according to claim 1 for preparing a compound of formula (1) which is 2-[4-(2,3-dihydro-1,4-benzodioxin-2-yl)methylamino]butyl-1,2-benzoisothiazol-3(2H)-one-1,1-dioxide or the
5 pharmaceutically acceptable salts thereof.

4. A process according to claim 1 for preparing a compound of formula (1) which is 2-[2-(2,3-dihydro-1,4-
10 -benzodioxin-2-yl)methylamino]ethyl-1,2-benzoisothiazol-3(2H)one-1,1-dioxide and the pharmaceutically acceptable salts thereof.

15 5. A process according to claim 1 for preparing a compound of formula (1) wherein X is

20



25


from 2 to 4, and the pharmaceutically acceptable salts thereof.

30

6. A process according to claim 1 for preparing a compound of formula (1) which is 2-[4-(2,3-dihydronaphtho-
[1,2-b[1,4]dioxin-3-yl)methylamino]butyl-1,2-benzoisothiazol-3(2H)one-1,1-dioxide and the pharma-
ceutically acceptable salts thereof.

35

7. A process according to claim 1 for preparing a compound of formula (1) wherein X is



5

n is an integer of from 2 to 4, and the pharmaceutically acceptable salts thereof.

10

8. A process according to claim 1 for preparing a compound of formula (1) which is 2-[4-(1,2,3,4-tetrahydro- β -carboliny1)butyl]-1,2-benzisothiazol-3-(2H)-one-1,1-dioxide and the pharmaceutically acceptable salts thereof.

9. Use of a compound of claim 1 to 8 for the preparation of a medicament for use as an anxiolytic agent.

15

10. Use of a compound of claim 1 to 8 for the preparation of a medicament for use as an antihypertensive agent.



EP 87 10 3077

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.4)
A	EP-A-0 060 610 (PFIZER INC.) * page 22, line 17 - page 23, line 7; page 30, entry 2; example 51 *	1,7,9- 11,13	C 07 D 275/06 C 07 D 417/12 C 07 D 471/04 A 61 K 31/425 A 61 K 31/535 A 61 K 31/54 A 61 K 31/40
A	EP-A-0 129 128 (TROPONWERKE GMBH & CO. KG) * claims 1-9 *	1,9-11 ,13	A 61 K 31/495 A 61 K 31/435// (C 07 D 471/04 C 07 D 221:00 C 07 D 209:00)
A	DE-A-2 856 528 (E.R. SQUIBB & SONS, INC.) * claims 1, 12, 15, 16 *	1,10, 13	
A	US-A-4 110 449 (P.C. WADE et al.) * claims 1, 18-20 *	1,10, 13	
A	US-A-2 751 392 (C.H. GROGAN et al.)		TECHNICAL FIELDS SEARCHED (Int. Cl.4)
A	US-A-4 405 622 (A.F. KLUGE)		C 07 D 275/00 C 07 D 417/00 C 07 D 491/00 A 61 K 31/00
The present search report has been drawn up for all claims			
Place of search BERLIN		Date of completion of the search 15-06-1987	Examiner VAN AMSTERDAM L.J.P.
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	